arXiv:1202.5740v1 [math.CA] 26 Feb 2012

Some estimates for commutators of fractional integrals associated to operators with Gaussian kernel bounds on weighted Morrey spaces

Hua Wang [*]

Department of Mathematics, Zhejiang University, Hangzhou 310027, China

Abstract

Let L be the infinitesimal generator of an analytic semigroup on $L^2(\mathbb{R}^n)$ with Gaussian kernel bound, and let $L^{-\alpha/2}$ be the fractional integrals of L for $0<\alpha<n$. In this paper, we will obtain some boundedness properties of commutators $\big[b, L^{-\alpha/2}\big]$ on the weighted Morrey spaces $L^{p,\kappa}(w)$ when the symbol b belongs to $BMO(\mathbb{R}^n)$ or homogeneous Lipschitz space.

MSC(2010): 42B20; 42B35

Keywords: Gaussian upper bound; fractional integrals; weighted Morrey spaces; commutators.

1 Introduction and main results

Suppose that L is the infinitesimal generator of an analytic semigroup $\{e^{-tL}\}_{t>0}$ on $L^2(\mathbb{R}^n)$ with a kernel $p_t(x,y)$ satisfying a Gaussian upper bound; that is, there exist positive constants C and A such that for all x, $y\in\mathbb{R}^n$ and all $t>0$, we have

$$|p_t(x,y)| \le \frac{C}{t^{n/2}} e^{-A\frac{|x-y|^2}{t}}. \tag{1.1}$$

Throughout this paper, we will always assume that the semigroup $\{e^{-tL}\}_{t>0}$ has a kernel which satisfies (1.1). This property is satisfied by a large class of differential operators, some examples can be seen in [7].

For any $0<\alpha<n$, the fractional integrals $L^{-\alpha/2}$ associated to the operator L is defined by

$$L^{-\alpha/2}f(x) = \frac{1}{\Gamma(\alpha/2)}\int_0^\infty e^{-tL}(f)(x)t^{\alpha/2-1}\,dt. \tag{1.2}$$

---

[*] E-mail address: wanghua@pku.edu.cn. Supported by National Natural Science Foundation of China under Grant #10871173 and #10931001.

Note that if $L = -\Delta$ is the Laplacian on $\mathbb{R}^n$, then $L^{-\alpha/2}$ is the classical fractional integral operator I_α, which is given by(see [20])

$$I_\alpha f(x) = \frac{\Gamma(\frac{n-\alpha}{2})}{2^\alpha \pi^{\frac{n}{2}} \Gamma(\frac{\alpha}{2})} \int_{\mathbb{R}^n} \frac{f(y)}{|x-y|^{n-\alpha}}\, dy.$$

Let b be a locally integrable function on $\mathbb{R}^n$. The commutator of b and $L^{-\alpha/2}$ is defined as follows

$$\big[b, L^{-\alpha/2}\big](f)(x) = b(x)L^{-\alpha/2}(f)(x) - L^{-\alpha/2}(bf)(x). \tag{1.3}$$

The first result on the theory of commutators was obtained by Coifman, Rochberg and Weiss in [3]. Since then, many authors have been interested in studying this theory. When $0 < \alpha < n$, $1 < p < n/\alpha$ and $1/q = 1/p - \alpha/n$, Chanillo [2] proved that the commutator $[b, I_\alpha]$ is bounded from $L^p(\mathbb{R}^n)$ to $L^q(\mathbb{R}^n)$ whenever $b \in BMO(\mathbb{R}^n)$. Paluszyński [18] showed that $b \in \dot{\Lambda}_\beta(\mathbb{R}^n)$ (homogeneous Lipschitz space) if and only if $[b, I_\alpha]$ is bounded from $L^p(\mathbb{R}^n)$ to $L^s(\mathbb{R}^n)$, where $0 < \beta < 1$, $1 < p < n/(\alpha+\beta)$ and $1/s = 1/p - (\alpha+\beta)/n$. For the weighted case, Segovia and Torrea [19] proved that when $b \in BMO(\mathbb{R}^n)$ and $w \in A_{p,q}$(Muckenhoupt weight class), then $[b, I_\alpha]$ is bounded from $L^p(w^p)$ to $L^q(w^q)$.

In 2004, by using a new sharp maximal function introduced in [14], Duong and Yan [7] extended the result of [2] from $(-\Delta)$ to the more general operator L defined above. More precisely, they showed that

Theorem A. *Let $0 < \alpha < n$, $1 < p < n/\alpha$ and $1/q = 1/p - \alpha/n$. If $b \in BMO(\mathbb{R}^n)$, then the commutator $\big[b, L^{-\alpha/2}\big]$ is bounded from $L^p(\mathbb{R}^n)$ to $L^q(\mathbb{R}^n)$.*

In 2008, Auscher and Martell [1] considered the weighted case and obtained the following result (see also [4]).

Theorem B. *Let $0 < \alpha < n$, $1 < p < n/\alpha$, $1/q = 1/p - \alpha/n$ and $w \in A_{p,q}$. If $b \in BMO(\mathbb{R}^n)$, then the commutator $\big[b, L^{-\alpha/2}\big]$ is bounded from $L^p(w^p)$ to $L^q(w^q)$.*

On the other hand, in 2009, Komori and Shirai [13] first introduced the weighted Morrey spaces $L^{p,\kappa}(w)$ which could be viewed as an extension of weighted Lebesgue spaces, and investigated the boundedness of the Hardy-Littlewood maximal operator, singular integral operator and fractional integral operator on these weighted spaces. Moreover, they also proved the following theorem.

Theorem C. *Let $0 < \alpha < n$, $1 < p < n/\alpha$, $1/q = 1/p - \alpha/n$, $0 < \kappa < p/q$ and $w \in A_{p,q}$. If $b \in BMO(\mathbb{R}^n)$, then the commutator $[b, I_\alpha]$ is bounded from $L^{p,\kappa}(w^p, w^q)$ to $L^{q,\kappa q/p}(w^q)$.*

The purpose of this paper is to study the boundedness of $\big[b, L^{-\alpha/2}\big]$ on the weighted Morrey spaces $L^{p,\kappa}(w)$ when $b \in BMO(\mathbb{R}^n)$ or $b \in \dot{\Lambda}_\beta(\mathbb{R}^n)$. Our main results in the paper are formulated as follows.

Theorem 1.1. *Let* $0<\alpha<n$, $1<p<n/\alpha$, $1/q=1/p-\alpha/n$, $0<\kappa<p/q$ *and* $w\in A_{p,q}$. *Suppose that* $b\in BMO(\mathbb{R}^n)$, *then the commutator* $\big[b,L^{-\alpha/2}\big]$ *is bounded from* $L^{p,\kappa}(w^p,w^q)$ *to* $L^{q,\kappa q/p}(w^q)$.

Theorem 1.2. *Let* $0<\beta<1$, $0<\alpha+\beta<n$, $1<p<n/(\alpha+\beta)$, $1/s=1/p-(\alpha+\beta)/n$ *and* $w\in A_{p,s}$. *Suppose that* $b\in\dot{\Lambda}_\beta(\mathbb{R}^n)$ *and* $0<\kappa<p/s$, *then the commutator* $\big[b,L^{-\alpha/2}\big]$ *is bounded from* $L^{p,\kappa}(w^p,w^s)$ *to* $L^{s,\kappa s/p}(w^s)$.

Theorem 1.3. *Let* $0<\beta<1$, $0<\alpha+\beta<n$, $1/s=1-(\alpha+\beta)/n$ *and* $w\in A_{1,s}$. *Suppose that* $b\in\dot{\Lambda}_\beta(\mathbb{R}^n)$ *and* $0<\kappa<1/s$, *then the commutator* $\big[b,L^{-\alpha/2}\big]$ *is bounded from* $L^{1,\kappa}(w,w^s)$ *to* $WL^{s,\kappa s}(w^s)$.

Theorem 1.4. *Let* $0<\beta<1$, $0<\alpha+\beta<n$, $1<p<n/(\alpha+\beta)$, $1/s=1/p-(\alpha+\beta)/n$ *and* $w\in A_{p,s}$. *Suppose that* $b\in\dot{\Lambda}_\beta(\mathbb{R}^n)$ *and* $\kappa=p/s$, *then the commutator* $\big[b,L^{-\alpha/2}\big]$ *is bounded from* $L^{p,\kappa}(w^p,w^s)$ *to* BMO_L.

2 Notations and definitions

First let us recall some standard definitions and notations. The classical A_p weight theory was first introduced by Muckenhoupt in the study of weighted L^p boundedness of Hardy-Littlewood maximal functions in [16]. Let w be a nonnegative, locally integrable function defined on $\mathbb{R}^n$, $B=B(x_0,r_B)$ denotes the ball with the center x_0 and radius r_B. Given a ball B and $\lambda>0$, λB denotes the ball with the same center as B whose radius is λ times that of B, we also denote the Lebesgue measure of B by $|B|$ and the weighted measure of B by $w(B)$, where $w(B)=\int_B w(x)\,dx$. We say that $w\in A_p$, $1<p<\infty$, if

$$\left(\frac{1}{|B|}\int_B w(x)\,dx\right)\left(\frac{1}{|B|}\int_B w(x)^{-1/(p-1)}\,dx\right)^{p-1}\le C\quad\text{for every ball } B\subseteq\mathbb{R}^n,$$

where C is a positive constant which is independent of B.

For the case $p=1$, $w\in A_1$, if

$$\frac{1}{|B|}\int_B w(x)\,dx\le C\cdot\operatorname*{ess\,inf}_{x\in B}w(x)\quad\text{for every ball } B\subseteq\mathbb{R}^n.$$

For the case $p=\infty$, $w\in A_\infty$ if it satisfies the A_p condition for some $1<p<\infty$.

We also need another weight class $A_{p,q}$ introduced by Muckenhoupt and Wheeden in [17]. A weight function w belongs to $A_{p,q}$ for $1<p<q<\infty$ if there exists a constant $C>0$ such that

$$\left(\frac{1}{|B|}\int_B w(x)^q\,dx\right)^{1/q}\left(\frac{1}{|B|}\int_B w(x)^{-p'}\,dx\right)^{1/p'}\le C\quad\text{for every ball } B\subseteq\mathbb{R}^n,$$

where p' denotes the conjugate exponent of $p>1$; that is, $1/p+1/p'=1$.

When $p = 1$, w is in the class $A_{1,q}$ with $1 < q < \infty$ if there exists a constant $C > 0$ such that

$$\left(\frac{1}{|B|}\int_B w(x)^q\,dx\right)^{1/q}\left(\operatorname*{ess\,sup}_{x\in B}\frac{1}{w(x)}\right) \le C \quad \text{for every ball } B \subseteq \mathbb{R}^n.$$

A weight function w is said to belong to the reverse Hölder class RH_r if there exist two constants $r > 1$ and $C > 0$ such that the following reverse Hölder inequality holds

$$\left(\frac{1}{|B|}\int_B w(x)^r\,dx\right)^{1/r} \le C\left(\frac{1}{|B|}\int_B w(x)\,dx\right) \quad \text{for every ball } B \subseteq \mathbb{R}^n.$$

We give the following results that we will use in the sequel.

Lemma 2.1 ([10])**.** *Let $w \in A_p$ with $p \ge 1$. Then, for any ball B, there exists an absolute constant $C > 0$ such that*

$$w(2B) \le C\,w(B).$$

In general, for any $\lambda > 1$, we have

$$w(\lambda B) \le C \cdot \lambda^{np} w(B),$$

where C does not depend on B nor on λ.

Lemma 2.2 ([11])**.** *Let $w \in RH_r$ with $r > 1$. Then there exists a constant $C > 0$ such that*

$$\frac{w(E)}{w(B)} \le C\left(\frac{|E|}{|B|}\right)^{(r-1)/r}$$

for any measurable subset E of a ball B.

Next we shall give the definitions of some function spaces. A locally integrable function b is said to be in $BMO(\mathbb{R}^n)$ if

$$\|b\|_* = \sup_B \frac{1}{|B|}\int_B |b(x) - b_B|\,dx < \infty,$$

where $b_B = \frac{1}{|B|}\int_B b(y)\,dy$ and the supremum is taken over all balls B in $\mathbb{R}^n$.

Theorem 2.3 ([6, 12])**.** *Assume that $b \in BMO(\mathbb{R}^n)$. Then for any $1 \le p < \infty$, we have*

$$\sup_B\left(\frac{1}{|B|}\int_B \big|b(x) - b_B\big|^p\,dx\right)^{1/p} \le C\|b\|_*.$$

Let $0 < \beta \le 1$. The homogeneous Lipschitz space $\dot{\Lambda}_\beta(\mathbb{R}^n)$ is defined by

$$\dot{\Lambda}_\beta(\mathbb{R}^n) = \Big\{b : \|b\|_{\dot{\Lambda}_\beta} = \sup_{x,h\in\mathbb{R}^n, h\neq 0}\frac{|b(x+h) - b(x)|}{|h|^\beta} < \infty\Big\}.$$

Given a Muckenhoupt's weight function w on $\mathbb{R}^n$, for $1 \le p < \infty$, we denote by $L^p(w)$ the space of all functions f satisfying

$$\|f\|_{L^p(w)} = \left(\int_{\mathbb{R}^n} |f(x)|^p w(x)\,dx \right)^{1/p} < \infty.$$

When $p = \infty$, $L^\infty(w)$ will be taken to mean $L^\infty(\mathbb{R}^n)$, and

$$\|f\|_{L^\infty(w)} = \|f\|_{L^\infty} = \operatorname*{ess\,sup}_{x \in \mathbb{R}^n} |f(x)|.$$

In [13], Komori and Shirai first defined the weighted Morrey spaces and obtained some known results relevant to this paper. For the boundedness of some other operators on these spaces, we refer the readers to [21, 22].

Definition 2.4 ([13])**.** *Let $1 \le p < \infty$, $0 < \kappa < 1$ and w be a weight function. Then the weighted Morrey space is defined by*

$$L^{p,\kappa}(w) = \left\{ f \in L^p_{loc}(w) : \|f\|_{L^{p,\kappa}(w)} < \infty \right\},$$

where

$$\|f\|_{L^{p,\kappa}(w)} = \sup_B \left(\frac{1}{w(B)^\kappa} \int_B |f(x)|^p w(x)\,dx \right)^{1/p}$$

and the supremum is taken over all balls B in $\mathbb{R}^n$.

We also denote by $WL^{p,\kappa}(w)$ the weighted weak Morrey space of all locally integrable functions satisfying

$$\|f\|_{WL^{p,\kappa}(w)} = \sup_B \sup_{t>0} \frac{1}{w(B)^{\kappa/p}} t \cdot w\big(\left\{ x \in B : |f(x)| > t \right\}\big)^{1/p} < \infty.$$

In order to deal with the fractional order case, we need to consider the weighted Morrey space with two weights.

Definition 2.5 ([13])**.** *Let $1 \le p < \infty$ and $0 < \kappa < 1$. Then for two weights u and v, the weighted Morrey space is defined by*

$$L^{p,\kappa}(u,v) = \left\{ f \in L^p_{loc}(u) : \|f\|_{L^{p,\kappa}(u,v)} < \infty \right\},$$

where

$$\|f\|_{L^{p,\kappa}(u,v)} = \sup_B \left(\frac{1}{v(B)^\kappa} \int_B |f(x)|^p u(x)\,dx \right)^{1/p}.$$

Theorem D ([13])**.** *Let $0 < \alpha < n$, $1 < p < n/\alpha$, $1/q = 1/p - \alpha/n$ and $w \in A_{p,q}$. Then the operator I_α is bounded from $L^{p,\kappa}(w^p, w^q)$ to $L^{q,\kappa q/p}(w^q)$.*

Theorem E ([13])**.** *Let $p = 1$, $0 < \alpha < n$, $1/q = 1 - \alpha/n$ and $w \in A_{1,q}$. Then the operator I_α is bounded from $L^{1,\kappa}(w, w^q)$ to $WL^{q,\kappa q}(w^q)$.*

We are going to conclude this section by defining the function spaces BMO_L. Duong and Yan [8] introduced and developed a new function space BMO_L associated with an operator L. Assume that the kernel $p_t(x,y)$ of $\{e^{-tL}\}_{t>0}$ satisfies an upper bound

$$|p_t(x,y)| \le t^{-n/2} g\Big(\frac{|x-y|}{t^{1/2}}\Big),$$

for all $x, y \in \mathbb{R}^n$ and all $t > 0$. Here g is a positive, bounded, decreasing function satisfying

$$\lim_{r\to\infty} r^{n+\epsilon} g(r) = 0, \quad \text{for some } \epsilon > 0. \tag{2.1}$$

Let ϵ be the constant in (2.1) and $0 < \beta < \epsilon$. A function $f \in L^p_{loc}(\mathbb{R}^n)$ is said to be a function of type (p,β) if f satisfies

$$\Big(\int_{\mathbb{R}^n} \frac{|f(x)|^p}{(1+|x|)^{n+\beta}}\,dx\Big)^{1/p} \le c < \infty. \tag{2.2}$$

We denote by $\mathcal{M}_{(p,\beta)}$ the collection of all functions of type (p,β). If $f \in \mathcal{M}_{(p,\beta)}$, then the norm of f in $\mathcal{M}_{(p,\beta)}$ is defined by

$$\|f\|_{\mathcal{M}_{(p,\beta)}} = \inf\big\{c \ge 0 : (2.2) \text{ holds}\big\}.$$

It is easy to see that $\mathcal{M}_{(p,\beta)}$ is a Banach space under the norm $\|f\|_{\mathcal{M}_{(p,\beta)}} < \infty$. We set

$$\mathcal{M}_p = \bigcup_{\beta:0<\beta<\epsilon} \mathcal{M}_{(p,\beta)}.$$

For any $f \in L^p(\mathbb{R}^n)$, $1 \le p < \infty$, Martell [14] defined a kind of sharp maximal function $M_L^{\#} f$ associated with the semigroup $\{e^{-tL}\}_{t>0}$ by the expression

$$M_L^{\#} f(x) = \sup_{x\in B} \frac{1}{|B|} \int_B \big|f(y) - e^{-t_B L} f(y)\big|\,dy,$$

where $t_B = r_B^2$ and r_B is the radius of the ball B. Let $f \in \mathcal{M}_p$ with $1 < p < \infty$, then we say that $f \in BMO_L$ if the sharp maximal function $M_L^{\#} f \in L^\infty(\mathbb{R}^n)$, and we define $\|f\|_{BMO_L} = \big\|M_L^{\#} f\big\|_{L^\infty}$. For further details about the properties and applications of BMO_L spaces, we refer the readers to [5, 8, 9].

Throughout this article, we will use C to denote a positive constant, which is independent of the main parameters and not necessarily the same at each occurrence. By $A \sim B$, we mean that there exists a constant $C > 1$ such that $\frac{1}{C} \le \frac{A}{B} \le C$. Moreover, we denote the conjugate exponent of $q > 1$ by $q' = q/(q-1)$.

3 Proofs of Theorems 1.1, 1.2 and 1.3

Proof. Fix a ball $B = B(x_0, r_B) \subseteq \mathbb{R}^n$ and decompose $f = f_1 + f_2$, where $f_1 = f\chi_{2B}$, χ_{2B} denotes the characteristic function of $2B$. Since $\big[b, L^{-\alpha/2}\big]$ is a linear operator, then we have

$$\begin{aligned}
&\frac{1}{w^q(B)^{\kappa/p}}\bigg(\int_B\big|\big[b,L^{-\alpha/2}\big]f(x)\big|^q w^q(x)\,dx\bigg)^{1/q}\\
\leq&\frac{1}{w^q(B)^{\kappa/p}}\bigg(\int_B\big|\big[b,L^{-\alpha/2}\big]f_1(x)\big|^q w^q(x)\,dx\bigg)^{1/q}+\frac{1}{w^q(B)^{\kappa/p}}\bigg(\int_B\big|\big[b,L^{-\alpha/2}\big]f_2(x)\big|^q w^q(x)\,dx\bigg)^{1/q}\\
=&I_1+I_2.
\end{aligned}$$

For the term I_1, since $w\in A_{p,q}$, then we get $w^q\in A_{1+q/p'}$(see [17]). Hence, it follows from Theorem B and Lemma 2.1 that

$$\begin{aligned}
I_1&\leq\frac{1}{w^q(B)^{\kappa/p}}\big\|\big[b,L^{-\alpha/2}\big]f_1\big\|_{L^q(w^q)}\\
&\leq C\|b\|_*\cdot\frac{1}{w^q(B)^{\kappa/p}}\|f_1\|_{L^p(w^p)}\\
&=C\|b\|_*\cdot\frac{1}{w^q(B)^{\kappa/p}}\bigg(\int_{2B}|f(x)|^p w^p(x)\,dx\bigg)^{1/p}\\
&\leq C\|b\|_*\|f\|_{L^{p,\kappa}(w^p,w^q)}\cdot\frac{w^q(2B)^{\kappa/p}}{w^q(B)^{\kappa/p}}\\
&\leq C\|b\|_*\|f\|_{L^{p,\kappa}(w^p,w^q)}.
\end{aligned}\tag{3.1}$$

We now turn to deal with the term I_2. Denote the kernel of $L^{-\alpha/2}$ by $K_\alpha(x,y)$, then for any $x\in B$, we write

$$\begin{aligned}
\big|\big[b,L^{-\alpha/2}\big]f_2(x)\big|&=\bigg|\int_{(2B)^c}\big[b(x)-b(y)\big]K_\alpha(x,y)f(y)\,dy\bigg|\\
&\leq\big|b(x)-b_B\big|\cdot\int_{(2B)^c}|K_\alpha(x,y)||f(y)|\,dy\\
&\quad+\int_{(2B)^c}|b(y)-b_B||K_\alpha(x,y)||f(y)|\,dy\\
&=\text{I+II}.
\end{aligned}$$

Since the kernel of e^{-tL} is $p_t(x,y)$, then it follows immediately from (1.2) that (see [15])

$$K_\alpha(x,y)=\frac{1}{\Gamma(\alpha/2)}\int_0^\infty p_t(x,y)t^{\alpha/2-1}\,dt.\tag{3.2}$$

Thus, by using the Gaussian upper bound (1.1) and the expression (3.2), we can deduce (see [7] and [15])

$$\begin{aligned}
|K_\alpha(x,y)|&\leq\frac{1}{\Gamma(\alpha/2)}\int_0^\infty|p_t(x,y)|t^{\alpha/2-1}\,dt\\
&\leq C\int_0^\infty e^{-A\frac{|x-y|^2}{t}}t^{\alpha/2-n/2-1}\,dt\\
&\leq C\cdot\frac{1}{|x-y|^{n-\alpha}}.
\end{aligned}\tag{3.3}$$

So we have

$$\mathrm{I} \le \big|b(x)-b_B\big| \cdot \sum_{k=1}^{\infty} \frac{1}{|2^{k+1}B|^{1-\alpha/n}} \int_{2^{k+1}B} |f(y)|\,dy.$$

By using Hölder's inequality and the fact that $w \in A_{p,q}$, we can get

$$\begin{aligned}\int_{2^{k+1}B} |f(y)|\,dy &\le \bigg(\int_{2^{k+1}B} |f(y)|^p w^p(y)\,dy\bigg)^{1/p} \bigg(\int_{2^{k+1}B} w^{-p'}(y)\,dy\bigg)^{1/p'}\\ &\le C\|f\|_{L^{p,\kappa}(w^p,w^q)} \big|2^{k+1}B\big|^{1/q+1/p'} \cdot \frac{1}{w^q(2^{k+1}B)^{1/q-\kappa/p}}\\ &= C\|f\|_{L^{p,\kappa}(w^p,w^q)} \big|2^{k+1}B\big|^{1-\alpha/n} \cdot \frac{1}{w^q(2^{k+1}B)^{1/q-\kappa/p}}. \qquad (3.4)\end{aligned}$$

Hence

$$\begin{aligned}&\frac{1}{w^q(B)^{\kappa/p}} \bigg(\int_B \mathrm{I}^q\, w^q(x)\,dx\bigg)^{1/q}\\ &\le C\|f\|_{L^{p,\kappa}(w^p,w^q)} \frac{1}{w^q(B)^{\kappa/p}} \sum_{k=1}^{\infty} \frac{1}{w^q(2^{k+1}B)^{1/q-\kappa/p}} \cdot \bigg(\int_B |b(x)-b_B|^q w^q(x)\,dx\bigg)^{1/q}\\ &= C\|f\|_{L^{p,\kappa}(w^p,w^q)} \sum_{k=1}^{\infty} \frac{w^q(B)^{1/q-\kappa/p}}{w^q(2^{k+1}B)^{1/q-\kappa/p}} \cdot \bigg(\frac{1}{w^q(B)} \int_B |b(x)-b_B|^q w^q(x)\,dx\bigg)^{1/q}.\end{aligned}$$

We now claim that for any $1 < q < \infty$ and $v \in A_\infty$, the following inequality holds

$$\bigg(\frac{1}{v(B)} \int_B |b(x)-b_B|^q v(x)\,dx\bigg)^{1/q} \le C\|b\|_*. \qquad (3.5)$$

In fact, since $v \in A_\infty$, then we know that there exists $r > 1$ such that $v \in RH_r$. Thus, by Hölder's inequality and Theorem 2.3, we obtain

$$\begin{aligned}\bigg(\frac{1}{v(B)} \int_B |b(x)-b_B|^q v(x)\,dx\bigg)^{1/q} &\le \frac{1}{v(B)^{1/q}} \bigg(\int_B |b(x)-b_B|^{qr'}\,dx\bigg)^{1/(qr')} \bigg(\int_B v(x)^r\,dx\bigg)^{1/(qr)}\\ &\le C\bigg(\frac{1}{|B|} \int_B |b(x)-b_B|^{qr'}\,dx\bigg)^{1/(qr')}\\ &\le C\|b\|_*,\end{aligned}$$

which is our desired result. Note that $w^q \in A_{1+q/p'} \subset A_\infty$. In addition, we have $w^q \in RH_r$ with $r > 1$. Thus, by Lemma 2.2, we get

$$\frac{w^q(B)}{w^q(2^{k+1}B)} \le C\left(\frac{|B|}{|2^{k+1}B|}\right)^{(r-1)/r}. \qquad (3.6)$$

Consequently

$$\begin{aligned}\frac{1}{w^q(B)^{\kappa/p}} \bigg(\int_B \mathrm{I}^q\, w^q(x)\,dx\bigg)^{1/q} &\le C\|b\|_* \|f\|_{L^{p,\kappa}(w^p,w^q)} \sum_{k=1}^{\infty} \left(\frac{1}{2^{kn}}\right)^{(1-1/r)(1/q-\kappa/p)}\\ &\le C\|b\|_* \|f\|_{L^{p,\kappa}(w^p,w^q)}, \qquad (3.7)\end{aligned}$$

where the last series is convergent since $r > 1$ and $0 < \kappa < p/q$. On the other hand

$$\begin{aligned}
\mathrm{II} &\le \sum_{k=1}^{\infty}\int_{2^{k+1}B\backslash 2^kB}|b(y)-b_B||K_\alpha(x,y)||f(y)|\,dy\\
&\le \sum_{k=1}^{\infty}\int_{2^{k+1}B\backslash 2^kB}\big|b(y)-b_{2^{k+1}B}\big||K_\alpha(x,y)||f(y)|\,dy\\
&\quad+\sum_{k=1}^{\infty}\int_{2^{k+1}B\backslash 2^kB}\big|b_{2^{k+1}B}-b_B\big||K_\alpha(x,y)||f(y)|\,dy\\
&=\mathrm{III}+\mathrm{IV}.
\end{aligned}$$

To estimate III and IV, we observe that when $x \in B$, $y \in (2B)^c$, then $|y-x| \sim |y-x_0|$. Hence, it follows directly from the kernel estimate (3.3) that

$$\mathrm{III} \le C\sum_{k=1}^{\infty}\frac{1}{|2^{k+1}B|^{1-\alpha/n}}\int_{2^{k+1}B}\big|b(y)-b_{2^{k+1}B}\big||f(y)|\,dy.$$

An application of Hölder's inequality yields

$$\begin{aligned}
&\int_{2^{k+1}B}\big|b(y)-b_{2^{k+1}B}\big||f(y)|\,dy\\
&\le\bigg(\int_{2^{k+1}B}\big|b(y)-b_{2^{k+1}B}\big|^{p'}w^{-p'}(y)\,dy\bigg)^{1/p'}\bigg(\int_{2^{k+1}B}|f(y)|^pw^p(y)\,dy\bigg)^{1/p}\\
&\le\|f\|_{L^{p,\kappa}(w^p,w^q)}\cdot w^q\big(2^{k+1}B\big)^{\kappa/p}\bigg(\int_{2^{k+1}B}\big|b(y)-b_{2^{k+1}B}\big|^{p'}w^{-p'}(y)\,dy\bigg)^{1/p'}.
\end{aligned}$$

We set $v(y) = w^{-p'}(y)$, then we have $v \in A_{1+p'/q} \subset A_\infty$ because $w \in A_{p,q}$ (see [17]). By the previous estimate (3.5) and the fact that $w \in A_{p,q}$, we obtain

$$\begin{aligned}
\bigg(\int_{2^{k+1}B}\big|b(y)-b_{2^{k+1}B}\big|^{p'}v(y)\,dy\bigg)^{1/p'}&\le C\|b\|_*v\big(2^{k+1}B\big)^{1/p'}\\
&\le C\|b\|_*\cdot\frac{|2^{k+1}B|^{1/q+1/p'}}{w^q(2^{k+1}B)^{1/q}}.
\end{aligned}\tag{3.8}$$

Note that $1/q + 1/p' = 1 - \alpha/n$. Hence, by (3.6) and (3.8), we have

$$\begin{aligned}
\frac{1}{w^q(B)^{\kappa/p}}\bigg(\int_B \mathrm{III}^q\,w^q(x)\,dx\bigg)^{1/q}&\le C\|b\|_*\|f\|_{L^{p,\kappa}(w^p,w^q)}\sum_{k=1}^{\infty}\frac{w^q(B)^{1/q-\kappa/p}}{w^q(2^{k+1}B)^{1/q-\kappa/p}}\\
&\le C\|b\|_*\|f\|_{L^{p,\kappa}(w^p,w^q)}.
\end{aligned}\tag{3.9}$$

Since $b \in BMO(\mathbb{R}^n)$, then a simple calculation gives that

$$\big|b_{2^{k+1}B}-b_B\big| \le C\cdot k\|b\|_*.$$

Thus, by the estimates (3.3) and (3.4), we get

$$\begin{aligned}
\mathrm{IV} &\le C\|b\|_* \sum_{k=1}^{\infty} k \cdot \frac{1}{|2^{k+1}B|^{1-\alpha/n}} \int_{2^{k+1}B} |f(y)|\,dy \\
&\le C\|b\|_*\|f\|_{L^{p,\kappa}(w^p,w^q)} \sum_{k=1}^{\infty} k \cdot \frac{1}{w^q(2^{k+1}B)^{1/q-\kappa/p}}.
\end{aligned}$$

Therefore

$$\begin{aligned}
\frac{1}{w^q(B)^{\kappa/p}}\left(\int_B \mathrm{IV}^q\, w^q(x)\,dx\right)^{1/q} &\le C\|b\|_*\|f\|_{L^{p,\kappa}(w^p,w^q)} \sum_{k=1}^{\infty} k \cdot \frac{w^q(B)^{1/q-\kappa/p}}{w^q(2^{k+1}B)^{1/q-\kappa/p}} \\
&\le C\|b\|_*\|f\|_{L^{p,\kappa}(w^p,w^q)} \sum_{k=1}^{\infty} \frac{k}{2^{kn\delta}} \\
&\le C\|b\|_*\|f\|_{L^{p,\kappa}(w^p,w^q)}, \qquad (3.10)
\end{aligned}$$

where $w^q \in RH_r$ and $\delta = (1-1/r)(1/q-\kappa/p)$. Summarizing the estimates (3.9) and (3.10) derived above, we thus obtain

$$\frac{1}{w^q(B)^{\kappa/p}}\left(\int_B \mathrm{II}^q\, w^q(x)\,dx\right)^{1/q} \le C\|b\|_*\|f\|_{L^{p,\kappa}(w^p,w^q)}. \qquad (3.11)$$

Combining the inequalities (3.1) and (3.7) with the above inequality (3.11) and taking the supremum over all balls $B \subseteq \mathbb{R}^n$, we complete the proof of Theorem 1.1. □

Obviously, by (3.3), we have the following pointwise inequality

$$\left|L^{-\alpha/2}(f)(x)\right| \le C \cdot I_\alpha(|f|)(x) \quad \text{for all } x \in \mathbb{R}^n.$$

Furthermore, by the definition of $b \in \dot{\Lambda}_\beta(\mathbb{R}^n)$ and (3.3), we deduce

$$\begin{aligned}
\left|\left[b, L^{-\alpha/2}\right](f)(x)\right| &\le \int_{\mathbb{R}^n} |b(x)-b(y)||K_\alpha(x,y)||f(y)|\,dy \\
&\le C\|b\|_{\dot{\Lambda}_\beta} \int_{\mathbb{R}^n} \frac{|f(y)|}{|x-y|^{n-\alpha-\beta}}\,dy \\
&\le C\|b\|_{\dot{\Lambda}_\beta} I_{\alpha+\beta}(|f|)(x). \qquad (3.12)
\end{aligned}$$

Hence, Theorems 1.2 and 1.3 follows immediately from Theorems D and E.

4 Proof of Theorem 1.4

Proof. For any given $x \in \mathbb{R}^n$, fix a ball $B = B(x_0, r_B)$ which contains x. We decompose $f = f_1 + f_2$, where $f_1 = f\chi_{2B}$, and set $t_B = r_B^2$. Then we write

$$\begin{aligned}
&\frac{1}{|B|}\int_B \Big|\big[b, L^{-\alpha/2}\big]f(y) - e^{-t_B L}\big[b, L^{-\alpha/2}\big]f(y)\Big|\, dy \\
&\le \frac{1}{|B|}\int_B \Big|\big[b, L^{-\alpha/2}\big]f_1(y)\Big|\, dy + \frac{1}{|B|}\int_B \Big|e^{-t_B L}\big[b, L^{-\alpha/2}\big]f_1(y)\Big|\, dy \\
&\quad + \frac{1}{|B|}\int_B \Big|\big[b, L^{-\alpha/2}\big]f_2(y) - e^{-t_B L}\big[b, L^{-\alpha/2}\big]f_2(y)\Big|\, dy \\
&= J_1 + J_2 + J_3.
\end{aligned}$$

We are now going to estimate each term respectively. For the first term J_1, since $w \in A_{p,s}$, then the operator $I_{\alpha+\beta}$ is bounded from $L^p(w^p)$ into $L^s(w^s)$(see [17]). We also know that $w^s \in A_{1+s/p'} \subset A_s$. Applying Hölder's inequality, the inequality (3.12), Lemma 2.1 and the fact that $w^s \in A_s$, we obtain

$$\begin{aligned}
J_1 &\le \frac{1}{|B|}\bigg(\int_B \big|\big[b, L^{-\alpha/2}\big]f_1(y)\big|^s w^s(y)\, dy\bigg)^{1/s}\bigg(\int_B w^{-s'}(y)\, dy\bigg)^{1/s'} \\
&\le C\|b\|_* \cdot \frac{1}{|B|}\bigg(\int_{2B} |f(y)|^p w^p(y)\, dy\bigg)^{1/p}\bigg(\int_B w^{-s'}(y)\, dy\bigg)^{1/s'} \\
&\le C\|b\|_*\|f\|_{L^{p,\kappa}(w^p,w^s)} \cdot \frac{w^s(2B)^{\kappa/p}}{w^s(B)^{1/s}} \\
&\le C\|b\|_*\|f\|_{L^{p,\kappa}(w^p,w^s)},
\end{aligned}$$

where the last inequality is due to our assumption $\kappa = p/s$. For the term J_2, since the kernel of $e^{-t_B L}$ is $p_{t_B}(y,z)$, then we may write

$$\begin{aligned}
J_2 &\le \frac{1}{|B|}\int_B \int_{\mathbb{R}^n} \big|p_{t_B}(y,z)\big|\big|\big[b, L^{-\alpha/2}\big]f_1(z)\big| dz dy \\
&\le \frac{1}{|B|}\int_B \int_{2B} \big|p_{t_B}(y,z)\big|\big|\big[b, L^{-\alpha/2}\big]f_1(z)\big| dz dy \\
&\quad + \sum_{k=1}^{\infty}\frac{1}{|B|}\int_B \int_{2^{k+1}B\backslash 2^k B} \big|p_{t_B}(y,z)\big|\big|\big[b, L^{-\alpha/2}\big]f_1(z)\big| dz dy \\
&= J_2' + J_2''.
\end{aligned}$$

For any $y \in B$ and $z \in 2B$, by (1.1), we have $\big|p_{t_B}(y,z)\big| \le C \cdot (t_B)^{-n/2}$. Thus

$$\begin{aligned}
J_2' &\le C \cdot \frac{1}{|B|}\int_B \int_{2B} \frac{1}{(t_B)^{n/2}}\big|\big[b, L^{-\alpha/2}\big]f_1(z)\big| dz dy \\
&\le C \cdot \frac{1}{|2B|}\int_{2B} \big|\big[b, L^{-\alpha/2}\big]f_1(z)\big| dz.
\end{aligned}$$

Using the same arguments as in the estimate of J_1, we can also deduce

$$J_2' \le C\|b\|_*\|f\|_{L^{p,\kappa}(w^p,w^s)}.$$

As before, we note that for any $y \in B$, $z \in (2B)^c$, then $|z-y| \sim |z-x_0|$. In this case, by using (1.1) again, we get $\big|p_{t_B}(y,z)\big| \le C \cdot \frac{(t_B)^{n/2}}{|y-z|^{2n}}$. Hence

$$\begin{aligned} J_2'' &\le C\sum_{k=1}^{\infty}\frac{1}{|B|}\int_B\int_{2^{k+1}B\backslash 2^kB}\frac{(t_B)^{n/2}}{|y-z|^{2n}}\big|\big[b,L^{-\alpha/2}\big]f_1(z)\big|dzdy\\ &\le C\sum_{k=1}^{\infty}\frac{1}{2^{kn}}\frac{1}{|2^{k+1}B|}\int_{2^{k+1}B}\big|\big[b,L^{-\alpha/2}\big]f_1(z)\big|dz. \end{aligned}$$

Following along the same lines as before, we can also show that

$$\begin{aligned} \int_{2^{k+1}B}\big|\big[b,L^{-\alpha/2}\big]f_1(z)\big|dz &\le C\bigg(\int_{2B}|f(z)|^pw^p(z)\,dz\bigg)^{1/p}\bigg(\int_{2^{k+1}B}w^{-s'}(z)\,dz\bigg)^{1/s'}\\ &\le C\|b\|_*\|f\|_{L^{p,\kappa}(w^p,w^s)}\frac{w^s(2B)^{\kappa/p}}{w^s(2^{k+1}B)^{1/s}}\cdot\big|2^{k+1}B\big|. \end{aligned}$$

Consequently

$$J_2'' \le C\|b\|_*\|f\|_{L^{p,\kappa}(w^p,w^s)}\sum_{k=1}^{\infty}\frac{1}{2^{kn}}\cdot\left(\frac{w^s(2B)}{w^s(2^{k+1}B)}\right)^{1/s}. \tag{4.1}$$

Observe that $w^s \in A_{1+s/p'}$, then there exists a number $r^* > 1$ such that $w^s \in RH_{r^*}$. Moreover, by using Lemma 2.2 again, we get

$$\frac{w^s(2B)}{w^s(2^{k+1}B)} \le C\left(\frac{|2B|}{|2^{k+1}B|}\right)^{(r^*-1)/r^*}. \tag{4.2}$$

Substituting the above inequality (4.2) into (4.1), we thus obtain

$$\begin{aligned} J_2'' &\le C\|b\|_*\|f\|_{L^{p,\kappa}(w^p,w^s)}\sum_{k=1}^{\infty}\left(\frac{1}{2^{kn}}\right)^{1+(r^*-1)/(sr^*)}\\ &\le C\|b\|_*\|f\|_{L^{p,\kappa}(w^p,w^s)}. \end{aligned}$$

Summarizing the estimates of J_2' and J_2'' derived above, we can get

$$J_2 \le C\|b\|_*\|f\|_{L^{p,\kappa}(w^p,w^s)}.$$

In order to estimate the last term J_3, we need the following result given in [5] (see also [7]).

Lemma 4.1. *For $0 < \alpha < n$, the difference operator $(I - e^{-tL})L^{-\alpha/2}$ has an associated kernel $\widetilde{K}_{\alpha,t}(y,z)$ which satisfies the following estimate*

$$\big|\widetilde{K}_{\alpha,t}(y,z)\big| \le \frac{C}{|y-z|^{n-\alpha}}\frac{t}{|y-z|^2}. \tag{4.3}$$

Hence, by the above kernel estimate (4.3) and the definition of $b \in \dot{\Lambda}_\beta(\mathbb{R}^n)$, we have

$$\begin{aligned}
J_3 &= \frac{1}{|B|}\int_B \Big|\big(I - e^{-t_B L}\big)L^{-\alpha/2}\big([b(y)-b(\cdot)]f_2\big)(y)\Big|\,dy \\
&\le \frac{1}{|B|}\int_B \int_{(2B)^c} \big|\widetilde{K}_{\alpha,t_B}(y,z)\big|\big|b(y)-b(z)\big|\big|f(z)\big|\,dz \\
&\le C\|b\|_{\dot{\Lambda}_\beta}\cdot\frac{1}{|B|}\int_B\int_{(2B)^c}\frac{1}{|y-z|^{n-\alpha-\beta}}\frac{r_B^2}{|y-z|^2}\big|f(z)\big|\,dzdy \\
&\le C\|b\|_{\dot{\Lambda}_\beta}\sum_{k=1}^\infty\frac{1}{2^{2k}}\frac{1}{|2^{k+1}B|^{1-(\alpha+\beta)/n}}\int_{2^{k+1}B}\big|f(z)\big|\,dz.
\end{aligned}$$

Since $w \in A_{p,s}$, then by using the estimate (3.4) and the fact that $\kappa = p/s$, we thus obtain

$$\begin{aligned}
\int_{2^{k+1}B}\big|f(z)\big|\,dz &\le C\|f\|_{L^{p,\kappa}(w^p,w^s)}\big|2^{k+1}B\big|^{1/s+1/p'}\cdot\frac{1}{w^s(2^{k+1}B)^{1/s-\kappa/p}} \\
&= C\|f\|_{L^{p,\kappa}(w^p,w^s)}\big|2^{k+1}B\big|^{1-(\alpha+\beta)/n}.
\end{aligned}$$

Therefore

$$\begin{aligned}
J_3 &\le C\|b\|_{\dot{\Lambda}_\beta}\|f\|_{L^{p,\kappa}(w^p,w^s)}\sum_{k=1}^\infty\frac{1}{2^{2k}} \\
&\le C\|b\|_{\dot{\Lambda}_\beta}\|f\|_{L^{p,\kappa}(w^p,w^s)}.
\end{aligned}$$

Combining the above estimates for J_1, J_2 and J_3 and taking the supremum over all balls $B \subseteq \mathbb{R}^n$, we finally conclude the proof of Theorem 1.4. ☐

References

[1] P. Auscher and J. M. Martell, Weighted norm inequalities for fractional operators, Indiana Univ. Math. J, **57**(2008), 1845–1870.

[2] S. Chanillo, A note on commutators, Indiana Univ. Math. J, **31**(1982), 7–16.

[3] R. R. Coifman, R. Rochberg and G. Weiss, Factorization theorems for Hardy spaces in several variables, Ann. of Math, **103**(1976), 611–635.

[4] D. Cruz-Uribe, J. M. Martell and C. Pérez, Extrapolation from A_∞ weights and applications, J. Funct. Anal, **213**(2004), 412–439.

[5] D. G. Deng, X. T. Duong, A. Sikora and L. X. Yan, Comparison of the classical BMO with the BMO spaces associated with operators and applications, Rev. Mat. Iberoamericana, **24**(2008), 267–296.

[6] J. Duoandikoetxea, Fourier Analysis, American Mathematical Society, Providence, Rhode Island, 2000.

[7] X. T. Duong and L. X. Yan, On commutators of fractional integrals, Proc. Amer. Math. Soc, **132**(2004), 3549–3557.

[8] X. T. Duong and L. X. Yan, New function spaces of BMO type, the John-Nirenberg inequality, interpolation and applications, Comm. Pure Appl. Math, **58**(2005), 1375–1420.

[9] X. T. Duong and L. X. Yan, Duality of Hardy and BMO spaces associated with operators with heat kernel bounds, J. Amer. Math. Soc, **18**(2005), 943–973.

[10] J. Garcia-Cuerva and J. Rubio de Francia, Weighted Norm Inequalities and Related Topics, North-Holland, Amsterdam, 1985.

[11] R. F. Gundy and R. L. Wheeden, Weighted integral inequalities for non-tangential maximal function, Lusin area integral, and Walsh-Paley series, Studia Math, **49**(1974), 107–124.

[12] F. John and L. Nirenberg, On functions of bounded mean oscillation, Comm. Pure Appl. Math, **14**(1961), 415–426.

[13] Y. Komori and S. Shirai, Weighted Morrey spaces and a singular integral operator, Math. Nachr, **282**(2009), 219–231.

[14] J. M. Martell, Sharp maximal functions associated with approximations of the identity in spaces of homogeneous type and applications, Studia Math, **161**(2004), 113–145.

[15] H. X. Mo and S. Z. Lu, Boundedness of multilinear commutators of generalized fractional integrals, Math. Nachr, **281**(2008), 1328–1340.

[16] B. Muckenhoupt, Weighted norm inequalities for the Hardy maximal function, Trans. Amer. Math. Soc, **165**(1972), 207–226.

[17] B. Muckenhoupt and R. L. Wheeden, Weighted norm inequalities for fractional integrals, Trans. Amer. Math. Soc, **192**(1974), 261–274.

[18] M. Paluszyński, Characterization of the Besov spaces via the commutator operator of Coifman, Rochberg and Weiss, Indiana Univ. Math. J, **44**(1995), 1–17.

[19] C. Segovia and J. L. Torrea, Weighted inequalities for commutators of fractional and singular integral, Publ. Mat, **35**(1991), 209–235.

[20] E. M. Stein, Singular Integrals and Differentiability Properties of Functions, Princeton Univ. Press, Princeton, New Jersey, 1970.

[21] H. Wang, The boundedness of some operators with rough kernel on the weighted Morrey spaces, Acta Math. Sinica(Chin. Ser), to appear.

[22] H. Wang and H. P. Liu, Some estimates for Bochner-Riesz operators on the weighted Morrey spaces, Acta Math. Sinica(Chin. Ser), to appear.